

# MS 221 — Homework Set (5)

---

## (Applications of The Chain Rule)

### QUESTION 1

A particle moving on a plane has Cartesian and polar coordinates at time  $t$  given by  $(x(t), y(t))$  and  $(r(t), \theta(t))$ , respectively. Thus,

$$x(t) = r(t) \cos \theta(t) \quad \text{and} \quad y(t) = r(t) \sin \theta(t) \quad \forall t \in \mathbf{R}.$$

If the speed in Cartesian coordinates is given by  $\sqrt{\dot{x}^2(t) + \dot{y}^2(t)}$  find the corresponding formula for the speed in terms of polar coordinates, that is, in terms of  $r, \theta, \dot{r}$  and  $\dot{\theta}$ .

### QUESTION 2

A disc with **centre at the origin** rotates anti-clockwise with **constant angular speed**  $\omega$  revolutions/sec about the origin. An insect on this disc is crawling in a straight line (relative to the disc) towards the centre at a constant speed (relative to the disc) of  $\alpha$  cm/sec. If the polar coordinates of the insect at time  $t = 0$  are  $r(0) = 100$  cm and  $\theta(0) = 0$  radians do the following:

- Find the polar coordinates  $(r(t), \theta(t))$  of the insect at any subsequent time  $t$ .
- Use part (a) to determine the Cartesian coordinates  $(x(t), y(t))$  of the insect at any subsequent time  $t$ .
- Find the velocity and acceleration (vectors) in Cartesian coordinates of the insect at any subsequent time  $t$ .

### QUESTION 3

A function  $f : \mathbf{R}^2 \rightarrow \mathbf{R} : (x, y) \mapsto f(x, y)$  satisfies

$$\frac{\partial f}{\partial x}(0, 0) = 3 \quad \text{and} \quad \frac{\partial f}{\partial y}(0, 0) = -5.$$

If in addition,  $f(ta, tb) = tf(a, b)$  for every  $t \in \mathbf{R}$  and for every  $(a, b) \in \mathbf{R}^2$ , find  $f(a, b) \forall (a, b) \in \mathbf{R}^2$ .

**Hint:**  $tf(a, b) \equiv f(ta, tb) \implies f(a, b) \equiv \frac{d}{dt}f(ta, tb).$

### QUESTION 4

Express the partial derivative  $\frac{\partial}{\partial x} f(u(x, y), v(x, y), w(x, y))$  in terms of the **Chain Rule**.

### QUESTION 5

Throughout this question  $\Omega$  will denote the set in the  $xy$ -plane given by:

$$\Omega = \{ (x, y) \in \mathbf{R}^2 \mid y > 0 \}.$$

If the functions  $\xi : \Omega \rightarrow \mathbf{R}$  and  $\eta : \Omega \rightarrow \mathbf{R}$  are specified by

$$\xi(x, y) = x \ln y \quad \text{and} \quad \eta(x, y) = x,$$

express the partial differential equation

$$x \frac{\partial u}{\partial x} - y \ln y \frac{\partial u}{\partial y} = u \quad \text{on } \Omega$$

as a partial differential equation in the  $(\xi, \eta)$  - coordinates and, hence or otherwise, solve this partial differential equation subject to the condition that

$$u(x, e) \equiv xe^x \quad \text{for all } x \in \mathbf{R}$$

### QUESTION 6

**Notation:** In the case where  $\omega = f(u(x, t), v(x, t))$  we will write  $\frac{\partial \omega}{\partial u} := \frac{\partial f}{\partial u}(u, v)$ ,

$$\frac{\partial \omega}{\partial v} := \frac{\partial f}{\partial v}(u, v), \quad \frac{\partial \omega}{\partial x} := \frac{\partial f}{\partial x}(u(x, t), v(x, t)), \quad \frac{\partial \omega}{\partial t} := \frac{\partial f}{\partial t}(u(x, t), v(x, t))$$

and similarly for higher order derivatives. Now, if

$$u(x, t) = x + ct \quad \text{and} \quad v(x, t) = x - ct,$$

where  $c$  is a non-zero constant, show that

$$\frac{\partial^2 \omega}{\partial x^2} - \frac{1}{c^2} \frac{\partial^2 \omega}{\partial t^2} \equiv 4 \frac{\partial^2 \omega}{\partial u \partial v}$$

### QUESTION 7

Find all solutions of the (partial differential) equation  $\frac{\partial^2 \omega}{\partial u \partial v} \equiv 0$ .

**Hint:** If a function  $h(r, s)$  satisfies  $\frac{\partial h}{\partial r} \equiv 0$ , then  $h$  is constant in  $r$ . That is,  $h$  is a function of  $s$  only.

### QUESTION 8

Use Questions 6 and 7 above to show that **every solution**  $\omega$  of the 1-dimensional wave equation

$$\frac{\partial^2 \omega}{\partial x^2} \equiv \frac{1}{c^2} \frac{\partial^2 \omega}{\partial t^2}$$

is of the form  $\omega = \varphi(x + ct) + \psi(x - ct)$  where  $\varphi$  and  $\psi$  are arbitrary smooth functions

# MS 221 HOMEWORK SET (5)

**Q1**

Write  $\dot{x}(t)$  for  $\frac{d\dot{x}}{dt}(t)$  etc.

$$x(t) = r(t) \cos \theta(t)$$

$$y(t) = r(t) \sin \theta(t)$$

$\Rightarrow$

$$\dot{x} = r \cos \theta - r \dot{\theta} \sin \theta$$

$$\dot{y} = r \sin \theta + r \dot{\theta} \cos \theta$$

$$\Rightarrow (\dot{x}^2 + \dot{y}^2) = (r \cos \theta - r \dot{\theta} \sin \theta)^2 + (r \sin \theta + r \dot{\theta} \cos \theta)^2$$

$$= \left( r^2 \cos^2 \theta - 2r r \dot{\theta} \cos \theta \sin \theta + r^2 \dot{\theta}^2 \sin^2 \theta \right)$$

$$+ \left( r^2 \sin^2 \theta + 2r r \dot{\theta} \cos \theta \sin \theta + r^2 \dot{\theta}^2 \cos^2 \theta \right)$$

$$= r^2 (\cos^2 \theta + \sin^2 \theta) + 0 + r^2 \dot{\theta}^2 (\sin^2 \theta + \cos^2 \theta)$$


$$= r^2 + r^2 \dot{\theta}^2$$

$$\Rightarrow \text{speed at time } t = \sqrt{\dot{x}^2 + \dot{y}^2}$$

$$= \sqrt{r^2 + r^2 \dot{\theta}^2}$$

L2

Q2



(a) Position of insect at time  $t$  in polar coordinates is  $(r(t), \theta(t))$   
where

$$r(t) = 100 - \alpha t \text{ cm.}$$

$$\theta(t) = 2\pi\omega t \text{ (radians)}$$

(b) From part (a)

$$x(t) = r(t) \cos \theta(t) = (100 - \alpha t) \cos 2\pi\omega t \text{ cm.}$$

$$y(t) = r(t) \sin \theta(t) = (100 - \alpha t) \sin 2\pi\omega t \text{ cm.}$$

(c) velocity =  $\begin{bmatrix} \dot{x} \\ \dot{y} \end{bmatrix} = \begin{bmatrix} -\alpha \cos 2\pi\omega t - 2\pi\omega(100 - \alpha t) \sin 2\pi\omega t \\ -\alpha \sin 2\pi\omega t + 2\pi\omega(100 - \alpha t) \cos 2\pi\omega t \end{bmatrix}$

acceleration =  $\begin{bmatrix} \ddot{x} \\ \ddot{y} \end{bmatrix} = \begin{bmatrix} 2\alpha(2\pi\omega)^2 \sin 2\pi\omega t - (2\pi\omega)^2(100 - \alpha t) \cos 2\pi\omega t \\ -2\alpha(2\pi\omega)^2 \cos 2\pi\omega t - (2\pi\omega)^2(100 - \alpha t) \sin 2\pi\omega t \end{bmatrix}$

3

Q3

$$t f(a, b) = f(ta, tb)$$

$$\Rightarrow f(a, b) = \frac{d}{dt} f(ta, tb)$$

$$= \frac{d}{dt} f(x(t), y(t)) \quad \begin{cases} x(t) = ta \\ y(t) = tb \end{cases}$$

$$= \frac{\partial f(p)}{\partial x} \frac{dx}{dt} + \frac{\partial f(p)}{\partial y} \frac{dy}{dt} \quad \begin{cases} \text{where} \\ p = (x(t), y(t)) \\ = (ta, tb) \end{cases}$$

In particular  
when  
 $t = 0$   
 $p = (0, 0)$

$$= \frac{\partial f(p)}{\partial x} \cdot a + \frac{\partial f(p)}{\partial y} \cdot b$$

$$= a \frac{\partial f}{\partial x}(0, 0) + b \frac{\partial f}{\partial y}(0, 0)$$

$$= 3a - 5b$$

Q4

$$\frac{\partial f(u(x, y), v(x, y), w(x, y))}{\partial x}$$

$$= \frac{\partial f(p)}{\partial u} \frac{\partial u}{\partial x}(x, y) + \frac{\partial f(p)}{\partial v} \frac{\partial v}{\partial x}(x, y) + \frac{\partial f(p)}{\partial w} \frac{\partial w}{\partial x}(x, y)$$

where  $p = (u(x, y), v(x, y), w(x, y))$

Q5

$$\xi(x, y) = x \ln y$$

$$\eta(x, y) = x$$

$$x \frac{\partial u}{\partial x} - y \ln y \frac{\partial u}{\partial y} = u \quad \text{on } \Omega$$

$$\Rightarrow x \left[ \frac{\partial u}{\partial \xi} \frac{\partial \xi}{\partial x} + \frac{\partial u}{\partial \eta} \frac{\partial \eta}{\partial x} \right] - y \ln y \left[ \frac{\partial u}{\partial \xi} \frac{\partial \xi}{\partial y} + \frac{\partial u}{\partial \eta} \frac{\partial \eta}{\partial y} \right] = u$$

$$\Rightarrow x \left[ \frac{\partial u}{\partial \xi} \cdot (\ln y) + \frac{\partial u}{\partial \eta} \cdot (1) \right] - y \ln y \left[ \cancel{\frac{\partial u}{\partial \xi} \cdot \left( \frac{x}{y} \right)} + \cancel{\frac{\partial u}{\partial \eta} \cdot (0)} \right] = u$$

$$\Rightarrow x \frac{\partial u}{\partial \eta} = u \quad \Rightarrow \boxed{\eta \frac{\partial u}{\partial \eta} = u}$$

$$\Rightarrow \int \frac{1}{u} \frac{\partial u}{\partial \eta} d\eta = \int \frac{1}{\eta} d\eta$$

$$\Rightarrow \int \frac{1}{u} du = \int \frac{1}{\eta} d\eta \quad (\text{integration by substitution})$$

$$\Rightarrow \ln u = \ln \eta + \ln \varphi(\xi) = \ln \eta \varphi(\xi)$$

$$\Rightarrow u(\xi, \eta) = \eta \varphi(\xi)$$

$$\Rightarrow u(x, y) := u(\xi(x, y), \eta(x, y)) = x \cdot \varphi(x \ln y)$$

$$\begin{cases} x e^x \equiv u(x, e) = x \varphi(x \cdot 1) \\ \Rightarrow \varphi(x) = e^x \end{cases}$$

$$\Rightarrow u(x, y) = x e^{x \ln y} = x e^{\ln y^x} = x y^x$$

Q6

$$u(x, t) = x + ct$$

$$v(x, t) = x - ct$$

5

$$\frac{\partial \omega}{\partial t} = \frac{\partial \omega}{\partial u} \frac{\partial u}{\partial t} + \frac{\partial \omega}{\partial v} \frac{\partial v}{\partial t}$$
$$\Rightarrow \omega_t = \omega_u \cdot (c) + \omega_v \cdot (-c)$$

write  
 $\omega_t = \frac{\partial \omega}{\partial t}$   
 $\omega_u = \frac{\partial \omega}{\partial u}$  etc

$$\Rightarrow \omega_t = c [\omega_u - \omega_v]$$

$$\Rightarrow \omega_{tt} = c \left[ \frac{\partial \omega_u}{\partial t} - \frac{\partial \omega_v}{\partial t} \right]$$

$$= c \left[ \frac{\partial \omega_u}{\partial u} \frac{\partial u}{\partial t} + \frac{\partial \omega_u}{\partial v} \frac{\partial v}{\partial t} - \left( \frac{\partial \omega_v}{\partial u} \frac{\partial u}{\partial t} + \frac{\partial \omega_v}{\partial v} \frac{\partial v}{\partial t} \right) \right]$$

$$= c \left[ \omega_{uu} \cdot (c) + \omega_{uv} \cdot (-c) - (\omega_{vu} \cdot (c) + \omega_{vv} \cdot (-c)) \right]$$

$$= c^2 [\omega_{uv} - 2\omega_{uv} + \omega_{vv}]$$

$$\Rightarrow \boxed{\omega_{tt} = c^2 [\omega_{uv} - 2\omega_{uv} + \omega_{vv}]} \quad \dots \textcircled{1}$$

Similarly

and

$$\omega_x = \omega_u + \omega_v$$

$$\boxed{\omega_{xx} = \omega_{uu} + 2\omega_{uv} + \omega_{vv}} \quad \dots \textcircled{2}$$

$$\Rightarrow \omega_{xx} - \frac{1}{c^2} \omega_{tt} = 4\omega_{uv} \quad \dots \textcircled{2} - \frac{1}{c^2} \textcircled{1}$$

16

Q 7


$$\frac{\partial^2 \omega}{\partial u \partial v} = 0$$

$$\text{i.e. } \frac{\partial}{\partial u} \left( \frac{\partial \omega}{\partial v} \right) = 0$$


$$\Rightarrow \frac{\partial \omega}{\partial v} = f(v)$$

$$\Rightarrow \omega = \int f_1(v) dv + f_2(u)$$

$$\Rightarrow \omega = \phi(v) + \psi(u)$$


  
 arbitrary smooth functions

L7



Q8

$$u = x + ct \quad \text{in Q5}$$

$$v = x - ct$$

and

$$4 \frac{\partial^2 \omega}{\partial u \partial v} = \frac{\partial^2 \omega}{\partial x^2} - \frac{1}{c^2} \frac{\partial^2 \omega}{\partial t^2} \stackrel{\text{wave eqn}}{=} 0$$

$$\Rightarrow 4 \frac{\partial^2 \omega}{\partial u \partial v} = 0$$

$$\Rightarrow \omega = \phi(u) + \psi(v) \quad \text{by Q6}$$

$$\Rightarrow \omega = \phi(x+ct) + \psi(x-ct)$$